Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

The Prevalence of Antibiotic-Resistant Staphylococcus aureus Nasal
Carriage among Industrial Hog Operation Workers, Community
Residents, and Children Living in Their Households: North
Carolina, USA

Sarah M. Hatcher, Sarah M. Rhodes, Jill R. Stewart, Ellen Silbergeld, Nora Pisanic, Jesper Larsen, Sharon Jiang, Amanda Krosche, Devon Hall, Karen C. Carroll, and Christopher D. Heaney

Table of Contents

Supplemental Methods

Assignment of *spa* types to clonal complexes (CCs)

Supplemental Tables

Table S1. Antibiotics and the concentrations used in antibiotic susceptibility testing

Table S2. Description of industrial hog operation (IHO) characteristics and IHO occupational activities reported by IHO worker participants, North Carolina, 2014

Table S3. *S. aureus spa*-types that occurred at least twice among adults and children (< 7 yr old) in the industrial hog operation (IHO) group and the community referent (CR) group, North Carolina, 2014

Table S4. *S. aureus spa* types for each participant with discordance in *spa* types in their two nasal swab *S. aureus* isolates, North Carolina, 2014

Additional Files

Supplemental Code and Data ZIP File

Excel File Table S1. Heat map of antimicrobial resistance, *scn* gene status, and *spa* type for *S. aureus* isolates for all participants by group.